

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.



Technical Memorandum 79642

Mineral Precipitation in North Slope Aufeis Fields

(NASA-TM-79642) MINERAL PRECIPITATION IN
NORTH SLOPE AUFEIS (NASA) 10 P HC A02/MF
A01 CSCL 08G

N79-10502

G3/43 Unclass
37155

Dorothy K. Hall

SEPTEMBER 1978

National Aeronautics and
Space Administration

Goddard Space Flight Center
Greenbelt, Maryland 20771



MINERAL PRECIPITATION IN NORTH SLOPE AUFEIS FIELDS

Dorothy K. Hall

Hydrospheric Sciences Branch

September 1978

GODDARD SPACE FLIGHT CENTER

Greenbelt, Maryland

MINERAL PRECIPITATION IN NORTH SLOPE AUFELS FIELDS

Dorothy K. Hall

Hydrospheric Sciences Branch

NASA/Goddard Space Flight Center

Greenbelt, Maryland

ABSTRACT

The Canning and Shaviovik River Aufeis Fields were studied on the ground and with aircraft data. Powdered calcium carbonate (CaCO_3) patches, a few cm in thickness, were found in discrete locations on both aufeis fields. This is indicative of chemical weathering of limestone bedrock which is known to underlie much of the eastern Arctic Coastal Plain of Alaska. Spring or river water which remains unfrozen throughout much of the winter carries CaCO_3 in solution; as the river ice freezes more deeply the CaCO_3 in solution is forced upwards through cracks in the river ice. Upon exposure to the cold air CaCO_3 is excluded as the water freezes, forming successive layers during aufeis growth. In the melt season CaCO_3 slush/powder accumulates in patches on top of the ice as the aufeis melts downward.

MINERAL PRECIPITATION IN NORTH SLOPE AUFELS FIELDS

Abundant evidence exists which documents the occurrence of mechanical weathering in the Arctic, but evidence for chemical weathering is much less apparent due to the lack of flowing water during much of the year and to cold temperatures which permit very low rates of degradation of organic matter to produce organic acids. Furthermore most weathering of rock material in cold regions has been thought to occur in the top few centimeters of the surface (Davies, 1969) as a result of freeze/thaw processes.

There is one type of weathering of which little mention has been made in the literature which contradicts some previously held beliefs about chemical weathering in Arctic regions. This is weathering of calcareous rock material by intrapermafrost spring water which remains unfrozen throughout the year. The surface manifestation is in the form of a white calcium carbonate (CaCO_3) powder/slush found on river icings or aufeis deposits. Aufeis is a unique type of ice which can form over an existing river ice cover in rivers underlain by impermeable material, in this case, permafrost. River or spring water is forced to the surface as river ice and the active layer coalesce upon freezing in a river bed.

In July, 1978 two melting aufeis fields on the Arctic Coastal Plain of Alaska were studied in the Canning and Shaviovik Rivers. A NASA Convair 990 aircraft was flown over the Arctic Coastal Plain of Alaska on July 12 for the purpose of studying various hydrologic features on the coastal plain using passive and active

microwave aircraft and satellite data. Measurements from air photos taken from the Convair 990 revealed that the Shaviovik Aufeis was 1.8 square km in area and the Canning Aufeis was 4.5 square km in area on July 12. On July 15 a field party studied the Shaviovik Aufeis (Figure 1) on the ground. Accumulations of a white powdery material were found on and within the ablating ice (Figure 2). This powder was often found in association with hummocks in the ice, presumably at the point where water under pressure emerges to the surface during aufeis formation. Using x-ray diffraction analysis this material was later determined to be pure calcium carbonate (CaCO_3) which was carried upward in solution in spring or river water. CaCO_3 and other mineral salts have been noted on other aufeis fields (Williams and Van Everdingen, 1975; Webber and Klein, 1977; C. S. Benson, written comm., 1978).

This CaCO_3 material was also found in discrete locations on the Canning Aufeis field. The accumulations of CaCO_3 were generally a few cm in thickness and caused irregular ablation of the ice in their vicinity. At least four discrete locations of this material were observed on the Canning Aufeis. This material was also observed in calving ice on the edge of the aufeis field 18 cm below the surface of the ice which was approximately 1 m in thickness.

Limestone bedrock is present in the Canning River area of the coastal plain (Hopkins et al., 1955). Spring and river water, high in CO_2 due to its low temperature, flows through the rocks eroding limestone as it flows; the high CO_2 content in the water facilitates solution of the limestone. CaCO_3 in solution

then flows to the surface under pressure. Water emanating from a spring freezes rapidly and the CaCO_3 and other material in solution is excluded during the freezing process. As the aufeis melts in the summer, this calcium carbonate is released into the river in the meltwater increasing the suspended sediment load in the water. Another mechanism for causing the presence of the CaCO_3 precipitate is downward freezing of river ice excluding impurities (of which CaCO_3 is one) as it freezes. These impurities become concentrated in the water remaining in the river channel (Benson, written comm., 1978). As this water is forced up under pressure, the water, with calcium carbonate in solution, freezes. If this is the case, then more CaCO_3 should occur in the top layers of aufeis because the top layers form last as successive overflows occur. As river ice continues to freeze downward the impurities become more concentrated in the water below.

Calcium carbonate is also significant in oceanography and glaciology. During aufeis melting, the CaCO_3 flows to the ocean during the summer providing a non-organic source of CaCO_3 to the Arctic Ocean. Dissolved calcium carbonate is of interest in chemical, biological and geochemical oceanography (Chen, 1978). The type of CaCO_3 discussed herein is unrelated to decomposition of organic matter which is the usual source of CaCO_3 in the oceans. This is another way in which CaCO_3 gets into the Arctic Ocean.

Calcium carbonate precipitate has been found beneath temperate glaciers (Hallett, 1976). CaCO_3 is dissolved from limestone bedrock in the regelation slip

process which is characteristic of the flow of temperate glaciers. Subglacial limestone deposits play a dominant role in modifying the glacial bed and may actually impede basal sliding according to Hallett (1976). This is significant because it shows that glacio-chemical processes are operating at the base of glaciers. Erosion of calcareous rock material under snow drifts has also been studied in Arctic regions (Davies, 1969).

In conclusion it appears that even in the absence of much organic acid available to dissolve limestone in the Arctic, substantial erosion of limestone occurs as in humid regions. Limestone in Arctic and desert regions composes highlands or cliffs whereas in humid regions it forms lowlands because it is easily eroded by rainwater and runoff containing organic acids. Furthermore it appears that weathering of calcareous rock material by chemical processes is occurring at depth and that weathering is not restricted to freeze/thaw processes operating close to the surface.

ACKNOWLEDGEMENTS

I would like to thank Mr. Imants Virsnieks of the University of Washington Polar Science Center, and Mr. Arne Hanson of the Naval Arctic Research Laboratory at Barrow, Alaska for invaluable field assistance, and personnel at the Naval Arctic Research Lab for logistical support. I would also like to thank Dr. Frank Carsey of NASA/Goddard Space Flight Center, for comments and suggestions regarding this paper, and Mr. Pete Garmiento, also of NASA, for performing the x-ray diffraction analysis.

REFERENCES

- Benson, C. S., 1978: Written Communication, August 1978.
- Chen, C. T. A., 1978: Decomposition of Calcium Carbonate and Organic Carbon in Deep Oceans. Science V. 201, pp. 735-736.
- Davis, J. L., 1969: Landforms of Cold Climates. M.I.T. Press, Cambridge, MA.
- Embleton, C. and C. A. M. King, 1975: Periglacial Geomorphology. John Wiley and Sons, NY
- Hallett, B., 1976: Deposits formed by Sublacial Precipitation of CaCO_3 . Geological Society of America Bulletin V. 87, pp. 1003-1015.
- Hopkins, D. M.; T. N. Karlstrom; R. F. Black; J. R. Williams; T. L. Péwé, A. T. Fernald and E. H. Muller, 1955: Permafrost and Ground Water in Alaska. Geological Survey Professional Paper 264-F.
- Webber, P. J. and D. R. Klein, 1977: Geobotanical and Ecological Observations at Two Locations in the West-Central Siberian Arctic. Arctic and Alpine Research V. 9, pp. 305-315.
- Williams, J. R. and R. O. Van Everdingan, 1975: Groundwater Investigations in Permafrost Regions of North America: A Review. North American Contribution to the Second International Permafrost Conference, pp. 435-446.




Figure 1. Air Photo of the Shaviovik River Aufeis Field on the Alaskan Arctic Coastal Plain, July 11, 1978.




Figure 2. Calcium Carbonate Slush on the Shaviovik River Aufeis Field, July 15, 1978.

ORIGINAL PAGE IS
OF POOR QUALITY